

## Functions

Slides by Christopher M. Bourke  
Instructor: Berthe Y. Choueiry

Fall 2007

Computer Science & Engineering 235  
Introduction to Discrete Mathematics  
Section 2.3 of Rosen  
[cse235@cse.unl.edu](mailto:cse235@cse.unl.edu)

## Introduction

You've already encountered *functions* throughout your education.

$$\begin{aligned}f(x, y) &= x + y \\f(x) &= x \\f(x) &= \sin x\end{aligned}$$

Here, however, we will study functions on *discrete* domains and ranges. Moreover, we generalize functions to mappings. Thus, there may not always be a "nice" way of writing functions like above.

## Definition

### Function

#### Definition

A *function*  $f$  from a set  $A$  to a set  $B$  is an assignment of exactly one element of  $B$  to *each element* of  $A$ . We write  $f(a) = b$  if  $b$  is the *unique* element of  $B$  assigned by the function  $f$  to the element  $a \in A$ . If  $f$  is a function from  $A$  to  $B$ , we write

$$f : A \rightarrow B$$

This can be read as " $f$  maps  $A$  to  $B$ ".

Note the subtlety:

- ▶ Each and every element in  $A$  has a *single* mapping.
- ▶ Each element in  $B$  *may* be mapped to by *several* elements in  $A$  or not at all.

## Definitions

### Terminology


#### Definition

Let  $f : A \rightarrow B$  and let  $f(a) = b$ . Then we use the following terminology:

- ▶  $A$  is the *domain* of  $f$ , denoted  $\text{dom}(f)$ .
- ▶  $B$  is the *codomain* of  $f$ .
- ▶  $b$  is the *image* of  $a$ .
- ▶  $a$  is the *preimage* (antecedent) of  $b$ .
- ▶ The *range* of  $f$  is the set of all images of elements of  $A$ , denoted  $\text{rng}(f)$ .

## Definitions

### Visualization



A function,  $f : A \rightarrow B$ .

## Definition I

### More Definitions

#### Definition

Let  $f_1$  and  $f_2$  be functions from a set  $A$  to  $\mathbb{R}$ . Then  $f_1 + f_2$  and  $f_1 f_2$  are also functions from  $A$  to  $\mathbb{R}$  defined by

$$\begin{aligned}(f_1 + f_2)(x) &= f_1(x) + f_2(x) \\(f_1 f_2)(x) &= f_1(x) f_2(x)\end{aligned}$$

#### Example

## Definition II

More Definitions

Let  $f_1(x) = x^4 + 2x^2 + 1$  and  $f_2(x) = 2 - x^2$  then

$$\begin{aligned}(f_1 + f_2)(x) &= (x^4 + 2x^2 + 1) + (2 - x^2) \\ &= x^4 + x^2 + 3 \\ (f_1 f_2)(x) &= (x^4 + 2x^2 + 1) \cdot (2 - x^2) \\ &= -x^6 + 3x^2 + 2\end{aligned}$$

### Definition

Let  $f : A \rightarrow B$  and let  $S \subseteq A$ . The *image* of  $S$  is the subset of  $B$  that consists of all the images of the elements of  $S$ . We denote the image of  $S$  by  $f(S)$ , so that

$$f(S) = \{f(s) \mid s \in S\}$$

## Definition III

More Definitions

Note that here, an *image* is a *set* rather than an element.

### Example

Let

- ▶  $A = \{a_1, a_2, a_3, a_4, a_5\}$
- ▶  $B = \{b_1, b_2, b_3, b_4\}$
- ▶  $f = \{(a_1, b_2), (a_2, b_3), (a_3, b_3), (a_4, b_1), (a_5, b_4)\}$
- ▶  $S = \{a_1, a_3\}$

Draw a diagram for  $f$ .

The *image* of  $S$  is  $f(S) = \{b_2, b_3\}$

### Definition

## Definition IV

More Definitions

A function  $f$  whose domain and codomain are subsets of the set of real numbers is called *strictly increasing* if  $f(x) < f(y)$  whenever  $x < y$  and  $x$  and  $y$  are in the domain of  $f$ . A function  $f$  is called *strictly decreasing* if  $f(x) > f(y)$  whenever  $x < y$  and  $x$  and  $y$  are in the domain of  $f$ .

## Injections, Surjections, Bijections I

Definitions

### Definition

A function  $f$  is said to be *one-to-one* (or *injective*) if

$$f(x) = f(y) \Rightarrow x = y$$

for all  $x$  and  $y$  in the domain of  $f$ . A function is an *injection* if it is one-to-one.

Intuitively, an injection simply means that each element in  $B$  has at most one preimage (antecedent).

It may be useful to think of the contrapositive of this definition:

$$x \neq y \Rightarrow f(x) \neq f(y)$$

## Injections, Surjections, Bijections II

Definitions

### Definition

A function  $f : A \rightarrow B$  is called *onto* (or *surjective*) if for every element  $b \in B$  there is an element  $a \in A$  with  $f(a) = b$ . A function is called a *surjection* if it is onto.

Again, intuitively, a surjection means that every element in the codomain is mapped. This implies that the range is the same as the codomain.

## Injections, Surjections, Bijections III

Definitions

### Definition


A function  $f$  is a *one-to-one correspondence* (or a *bijection*, if it is *both* one-to-one and onto.

One-to-one correspondences are important because they endow a function with an *inverse*. They also allow us to have a concept of cardinality for infinite sets!

Let's take a look at a few general examples to get the feel for these definitions.

## Function Examples


A Non-function



This is not a function: Both  $a_1$  and  $a_2$  map to more than one element in  $B$ .

## Function Examples


A Function; Neither One-To-One Nor Onto



This function not one-to-one since  $a_1$  and  $a_3$  both map to  $b_1$ . It is not onto either since  $b_4$  is not mapped to by any element in  $A$ .

## Function Examples


One-To-One, Not Onto



This function is one-to-one since every  $a_i \in A$  maps to a unique element in  $B$ . However, it is not onto since  $b_4$  is not mapped to by any element in  $A$ .

## Function Examples


Onto, Not One-To-One



This function is onto since every element  $b_i \in B$  is mapped to by some element in  $A$ . However, it is not one-to-one since  $b_3$  is mapped to more than one element in  $A$ .

## Function Examples

A Bijection



This function is a bijection because it is both one-to-one and onto; every element in  $A$  maps to a unique element in  $B$  and every element in  $B$  is mapped by some element in  $A$ .

## Exercises I

Exercise I

### Example

Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  be defined by

$$f(x) = 2x - 3$$

What is the domain and range of  $f$ ? Is it onto? One-to-one?

Clearly,  $\text{dom}(f) = \mathbb{Z}$ . To see what the range is, note that

$$\begin{aligned} b \in \text{rng}(f) &\iff b = 2a - 3 & a \in \mathbb{Z} \\ &\iff b = 2(a - 2) + 1 \\ &\iff b \text{ is odd} \end{aligned}$$

## Exercises II

### Exercise I

Therefore, the range is the set of all *odd* integers. Since the range and codomain are different, (i.e.  $\text{rng}(f) \neq \mathbb{Z}$ ) we can also conclude that  $f$  is *not* onto.

However,  $f$  is one-to-one. To prove this, note that

$$\begin{aligned} f(x_1) = f(x_2) &\Rightarrow 2x_1 - 3 = 2x_2 - 3 \\ &\Rightarrow x_1 = x_2 \end{aligned}$$

follows from simple algebra.

## Exercises

### Exercise II

#### Example

Let  $f$  be as before,

$$f(x) = 2x - 3$$

but now define  $f : \mathbb{N} \rightarrow \mathbb{N}$ . What is the domain and range of  $f$ ? Is it onto? One-to-one?

By changing the domain/codomain in this example,  $f$  is not even a function anymore. Consider  $f(1) = 2 \cdot 1 - 3 = -1 \notin \mathbb{N}$ .

## Exercises I

### Exercise III

#### Example

Define  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  by

$$f(x) = x^2 - 5x + 5$$

Is this function one-to-one? Onto?

It is not one-to-one since for

$$\begin{aligned} f(x_1) = f(x_2) &\Rightarrow x_1^2 - 5x_1 + 5 = x_2^2 - 5x_2 + 5 \\ &\Rightarrow x_1^2 - 5x_1 = x_2^2 - 5x_2 \\ &\Rightarrow x_1^2 - x_2^2 = 5x_1 - 5x_2 \\ &\Rightarrow (x_1 - x_2)(x_1 + x_2) = 5(x_1 - x_2) \\ &\Rightarrow (x_1 + x_2) = 5 \end{aligned}$$

## Exercises II

### Exercise III

Therefore, any  $x_1, x_2 \in \mathbb{Z}$  satisfies the equality (i.e. there are an infinite number of solutions). In particular  $f(2) = f(3) = -1$ .

It is also *not* onto. The function is a parabola with a global minimum (calculus exercise) at  $(\frac{5}{2}, -\frac{9}{4})$ . Therefore, the function fails to map to any integer less than  $-1$ .

What would happen if we changed the domain/codomain?

## Exercises I

### Exercise IV

#### Example

Define  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  by

$$f(x) = 2x^2 + 7x$$

Is this function one-to-one? Onto?

Again, since this is a parabola, it cannot be onto (where is the global minimum?).

## Exercises II

### Exercise IV

However, it *is* one-to-one. We follow a similar argument as before:

$$\begin{aligned} f(x_1) = f(x_2) &\Rightarrow 2x_1^2 + 7x_1 = 2x_2^2 + 7x_2 \\ &\Rightarrow 2(x_1 - x_2)(x_1 + x_2) = 7(x_2 - x_1) \\ &\Rightarrow (x_1 + x_2) = \frac{7}{2} \end{aligned}$$

But  $\frac{7}{2} \notin \mathbb{Z}$  therefore, it must be the case that  $x_1 = x_2$ . It follows that  $f$  is one-to-one.

## Exercises I

### Exercise V

#### Example

Define  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  by

$$f(x) = 3x^3 - x$$

Is  $f$  one-to-one? Onto?

To see if its one-to-one, again suppose that  $f(x_1) = f(x_2)$  for  $x_1, x_2 \in \mathbb{Z}$ . Then

$$\begin{aligned} 3x_1^3 - x_1 &= 3x_2^3 - x_2 \Rightarrow 3(x_1^3 - x_2^3) = (x_1 - x_2) \\ &\Rightarrow 3(x_1 - x_2)(x_1^2 + x_1x_2 + x_2^2) = (x_1 - x_2) \\ &\Rightarrow (x_1^2 + x_1x_2 + x_2^2) = \frac{1}{3} \end{aligned}$$

## Exercises II

### Exercise V

Again, this is impossible since  $x_1, x_2$  are integers, thus  $f$  is one-to-one.

However, the function is *not* onto. Consider this counter example:  $f(a) = 1$  for some integer  $a$ . If this were true, then it must be the case that

$$a(3a^2 - 1) = 1$$

Where  $a$  and  $(3a^2 - 1)$  are integers. But the only time we can ever get that the product of two integers is 1 is when we have  $-1(-1)$  or  $1(1)$  neither of which satisfy the equality.

## Inverse Functions I

#### Definition

Let  $f : A \rightarrow B$  be a bijection. The *inverse function* of  $f$  is the function that assigns to an element  $b \in B$  the unique element  $a \in A$  such that  $f(a) = b$ . The inverse function of  $f$  is denoted by  $f^{-1}$ . Thus  $f^{-1}(b) = a$  when  $f(a) = b$ .

More succinctly, if an inverse exists,

$$f(a) = b \iff f^{-1}(b) = a$$

## Inverse Functions II


Note that by the definition, a function can have an inverse if and only if it is a bijection. Thus, we say that a bijection is *invertible*.

Why must a function be bijective to have an inverse?

- ▶ Consider the case where  $f$  is not one-to-one. This means that some element  $b \in B$  is mapped to by more than one element in  $A$ ; say  $a_1$  and  $a_2$ . How can we define an inverse? Does  $f^{-1}(b) = a_1$  or  $a_2$ ?
- ▶ Consider the case where  $f$  is not onto. This means that there is some element  $b \in B$  that is not mapped to by any  $a \in A$ , therefore what is  $f^{-1}(b)$ ?

## Inverse Functions

### Figure



A function & its inverse.

## Examples

### Example I

#### Example

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = 2x - 3$$

What is  $f^{-1}$ ?

First, verify that  $f$  is a bijection (it is). To find an inverse, we use substitution:

- ▶ Let  $f^{-1}(y) = x$
- ▶ Let  $y = 2x - 3$  and solve for  $x$
- ▶ Clearly,  $x = \frac{y+3}{2}$  so,
- ▶  $f^{-1}(y) = \frac{y+3}{2}$ .

## Examples

### Example II

#### Example

Let

$$f(x) = x^2$$

What is  $f^{-1}$ ?

No domain/codomain has been specified. Say  $f : \mathbb{R} \rightarrow \mathbb{R}$  Is  $f$  a bijection? Does an inverse exist?

No, however if we specify that

$$A = \{x \in \mathbb{R} \mid x \leq 0\}$$

and

$$B = \{y \in \mathbb{R} \mid y \geq 0\}$$

then it becomes a bijection and thus has an inverse.

## Examples

### Example II Continued

To find the inverse, we again, let  $f^{-1}(y) = x$  and  $y = x^2$ . Solving for  $x$  we get  $x = \pm\sqrt{y}$ . But which is it?

Since  $\text{dom}(f)$  is all nonpositive and  $\text{rng}(f)$  is nonnegative,  $y$  must be positive, thus

$$f^{-1}(y) = -\sqrt{y}$$

Thus, it should be clear that domains/codomains are just as important to a function as the definition of the function itself.

## Examples

### Example III

#### Example

Let

$$f(x) = 2^x$$

What should the domain/codomain be for this to be a bijection? What is the inverse?

The function should be  $f : \mathbb{R} \rightarrow \mathbb{R}^+$ . What happens when we include 0? Restrict either one to  $\mathbb{Z}$ ?

Let  $f^{-1}(y) = x$  and  $y = 2^x$ , solving for  $x$  we get  $x = \log_2(x)$ .

Therefore,

$$f^{-1}(y) = \log_2(y)$$

## Composition I

The values of functions can be used as the input to other functions.

#### Definition

Let  $g : A \rightarrow B$  and let  $f : B \rightarrow C$ . The *composition* of the functions  $f$  and  $g$  is

$$(f \circ g)(x) = f(g(x))$$

## Composition II

Note the *order* that you apply a function matters—you go from inner most to outer most.


The composition  $f \circ g$  cannot be defined unless the the range of  $g$  is a subset of the domain of  $f$ ;

$$f \circ g \text{ is defined} \iff \text{rng}(g) \subseteq \text{dom}(f)$$

It also follows that  $f \circ g$  is not necessarily the same as  $g \circ f$ .

## Composition of Functions

Figure



The composition of two functions.

## Composition

### Example I

#### Example

Let  $f$  and  $g$  be functions,  $\mathbb{R} \rightarrow \mathbb{R}$  defined by

$$\begin{aligned}f(x) &= 2x - 3 \\g(x) &= x^2 + 1\end{aligned}$$

What are  $f \circ g$  and  $g \circ f$ ?

Note that  $f$  is bijective, thus  $\text{dom}(f) = \text{rng}(f) = \mathbb{R}$ . For  $g$ , we have that  $\text{dom}(g) = \mathbb{R}$  but that  $\text{rng}(g) = \{x \in \mathbb{R} \mid x \geq 1\}$ .

## Composition

### Example I

Even so,  $\text{rng}(g) \subseteq \text{dom}(f)$  and so  $f \circ g$  is defined. Also,  $\text{rng}(f) \subseteq \text{dom}(g)$  so  $g \circ f$  is defined as well.

$$\begin{aligned}(f \circ g)(x) &= g(f(x)) \\&= g(2x - 3) \\&= (2x - 3)^2 + 1 \\&= 4x^2 - 12x + 10\end{aligned}$$

and

$$\begin{aligned}(g \circ f)(x) &= f(g(x)) \\&= f(x^2 + 1) \\&= 2(x^2 + 1) - 3 \\&= 2x^2 - 1\end{aligned}$$

## Equality

Though intuitive, we formally state what it means for two functions to be equal.

#### Lemma

Two functions  $f$  and  $g$  are equal if and only if  $\text{dom}(f) = \text{dom}(g)$  and

$$\forall a \in \text{dom}(f)(f(a) = g(a))$$

## Associativity

Though the composition of functions is not commutative ( $f \circ g \neq g \circ f$ ), it *is associative*.

#### Lemma

Composition of functions is an associative operation; that is,

$$(f \circ g) \circ h = f \circ (g \circ h)$$

## Important Functions

### Identity Function

#### Definition

The *identity function* on a set  $A$  is the function

$$\iota : A \rightarrow A$$

defined by  $\iota(a) = a$  for all  $a \in A$ . This symbol is the Greek letter iota.

One can view the identity function as a composition of a function and its inverse;

$$\iota(a) = (f \circ f^{-1})(a)$$

Moreover, the composition of any function  $f$  with the identity function is itself  $f$ ;

$$(f \circ \iota)(a) = (\iota \circ f)(a) = f(a)$$

## Inverses & Identity

The identity function, along with the composition operation gives us another characterization for when a function has an inverse.

#### Theorem

Functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$  are inverses if and only if

$$g \circ f = \iota_A \text{ and } f \circ g = \iota_B$$

That is,

$$\forall a \in A, b \in B ((g(f(a)) = a \wedge f(g(b)) = b)$$

## Important Functions I

### Absolute Value Function

#### Definition

The *absolute value* function, denoted  $|x|$  is a function  $f: \mathbb{R} \rightarrow \{y \in \mathbb{R} \mid y \geq 0\}$ . Its value is defined by

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

## Floor & Ceiling Functions


#### Definition

The *floor function*, denoted  $\lfloor x \rfloor$  is a function  $\mathbb{R} \rightarrow \mathbb{Z}$ . Its value is the largest integer that is less than or equal to  $x$ .

The *ceiling function*, denoted  $\lceil x \rceil$  is a function  $\mathbb{R} \rightarrow \mathbb{Z}$ . Its value is the smallest integer that is greater than or equal to  $x$ .

## Floor & Ceiling Functions

### Graphical View



## Factorial Function

The factorial function gives us the number of permutations (that is, uniquely ordered arrangement) of a collection of  $n$  objects.

#### Definition

The *factorial function*, denoted  $n!$  is a function  $\mathbb{N} \rightarrow \mathbb{Z}^+$ . Its value is the product of the first  $n$  positive integers.

$$n! = \prod_{i=1}^n i = 1 \cdot 2 \cdot 3 \cdots (n-1) \cdot n$$

## Factorial Function

### Stirling's Approximation

The factorial function is defined on a discrete domain. In many applications, it is useful to consider a continuous version of the function (say if we want to differentiate it).

To this end, we have *Stirling's Formula*:

$$n! \approx \sqrt{2\pi n} \frac{n^n}{e^n}$$