

Predicate Logic and Quantifiers

Slides by Christopher M. Bourke
Instructor: Berthe Y. Choueiry

Fall 2007

Computer Science & Engineering 235
Introduction to Discrete Mathematics
Sections 1.3–1.4 of Rosen

cse235@cse.unl.edu

Consider the following statements:

$$x > 3, \quad x = y + 3, \quad x + y = z$$

The truth value of these statements has no meaning without specifying the values of x, y, z .

However, we *can* make propositions out of such statements.

A *predicate* is a property that is affirmed or denied about the *subject* (in logic, we say “variable” or “argument”) of a *statement*.

$$\text{“} \underbrace{x}_{\text{subject}} \text{ is greater than } \underbrace{3}_{\text{predicate}} \text{”}$$

Terminology: affirmed = holds = is true; denied = does not hold = is not true.

To write in predicate logic:

“ x is greater than 3 ”

subject predicate

We introduce a (functional) symbol for the predicate, and put the subject as an argument (to the functional symbol): $P(x)$

Examples:

- $\text{Father}(x)$: unary predicate
- $\text{Brother}(x,y)$: binary predicate
- $\text{Sum}(x,y,z)$: ternary predicate
- $P(x,y,z,t)$: n -ary predicate

Definition

A statement of the form $P(x_1, x_2, \dots, x_n)$ is the value of the *propositional function* P . Here, (x_1, x_2, \dots, x_n) is an n -tuple and P is a predicate.

You can think of a propositional function as a function that

- Evaluates to true or false.
- Takes one or more arguments.
- Expresses a predicate involving the argument(s).
- Becomes a proposition when values are assigned to the arguments.

Propositional Functions

Example

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Universe of
Discourse

Quantifiers

Logic
Programming

Transcribing
English into
Logic

Further
Examples &
Exercises

Example

Let $Q(x, y, z)$ denote the statement " $x^2 + y^2 = z^2$ ". What is the truth value of $Q(3, 4, 5)$? What is the truth value of $Q(2, 2, 3)$? How many values of (x, y, z) make the predicate true?

Propositional Functions

Example

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Universe of
Discourse

Quantifiers

Logic
Programming

Transcribing
English into
Logic

Further
Examples &
Exercises

Example

Let $Q(x, y, z)$ denote the statement " $x^2 + y^2 = z^2$ ". What is the truth value of $Q(3, 4, 5)$? What is the truth value of $Q(2, 2, 3)$? How many values of (x, y, z) make the predicate true?

Since $3^2 + 4^2 = 25 = 5^2$, $Q(3, 4, 5)$ is true.

Since $2^2 + 2^2 = 8 \neq 3^2 = 9$, $Q(2, 2, 3)$ is false.

There are infinitely many values for (x, y, z) that make this propositional function true—how many right triangles are there?

Consider the previous example. Does it make sense to assign to x the value “blue”?

Intuitively, the *universe of discourse* is the set of all things we wish to talk about; that is, the set of all objects that we can sensibly assign to a variable in a propositional function.

What would be the universe of discourse for the propositional function $P(x) =$ “The test will be on x the 23rd” be?

Moreover, each variable in an n -tuple may have a different universe of discourse.

Let $P(r, g, b, c) =$ “The rgb-value of the color c is (r, g, b) ”.

For example, $P(255, 0, 0, \text{red})$ is true, while $P(0, 0, 255, \text{green})$ is false.

What are the universes of discourse for (r, g, b, c) ?

A predicate becomes a proposition when we assign it fixed values. However, another way to make a predicate into a proposition is to *quantify* it. That is, the predicate is true (or false) for *all* possible values in the universe of discourse or for *some* value(s) in the universe of discourse.

Such *quantification* can be done with two *quantifiers*: the *universal* quantifier and the *existential* quantifier.

Definition

The *universal quantification* of a predicate $P(x)$ is the proposition “ $P(x)$ is true for all values of x in the universe of discourse” We use the notation

$$\forall x P(x)$$

which can be read “for all x ”

If the universe of discourse is finite, say $\{n_1, n_2, \dots, n_k\}$, then the universal quantifier is simply the conjunction of all elements:

$$\forall x P(x) \iff P(n_1) \wedge P(n_2) \wedge \dots \wedge P(n_k)$$

Universal Quantifier

Example I

- Let $P(x)$ be the predicate “ x must take a discrete mathematics course” and let $Q(x)$ be the predicate “ x is a computer science student”.
- The universe of discourse for both $P(x)$ and $Q(x)$ is all UNL students.
- Express the statement “Every computer science student must take a discrete mathematics course”.

- Express the statement “Everybody must take a discrete mathematics course or be a computer science student”.

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic

Programming

Transcribing
English into
Logic

Further

Universal Quantifier

Example I

- Let $P(x)$ be the predicate “ x must take a discrete mathematics course” and let $Q(x)$ be the predicate “ x is a computer science student”.
- The universe of discourse for both $P(x)$ and $Q(x)$ is all UNL students.
- Express the statement “Every computer science student must take a discrete mathematics course”.

$$\forall x(Q(x) \rightarrow P(x))$$

- Express the statement “Everybody must take a discrete mathematics course or be a computer science student”.

- Let $P(x)$ be the predicate “ x must take a discrete mathematics course” and let $Q(x)$ be the predicate “ x is a computer science student”.
- The universe of discourse for both $P(x)$ and $Q(x)$ is all UNL students.
- Express the statement “Every computer science student must take a discrete mathematics course”.

$$\forall x(Q(x) \rightarrow P(x))$$

- Express the statement “Everybody must take a discrete mathematics course or be a computer science student”.

$$\forall x(Q(x) \vee P(x))$$

Universal Quantifier

Example I

- Let $P(x)$ be the predicate “ x must take a discrete mathematics course” and let $Q(x)$ be the predicate “ x is a computer science student”.
- The universe of discourse for both $P(x)$ and $Q(x)$ is all UNL students.
- Express the statement “Every computer science student must take a discrete mathematics course”.

$$\forall x(Q(x) \rightarrow P(x))$$

- Express the statement “Everybody must take a discrete mathematics course or be a computer science student”.

$$\forall x(Q(x) \vee P(x))$$

- Are these statements true or false?

Universal Quantifier

Example II

Express the statement “for every x and for every y , $x + y > 10$ ”

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Universal Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “for every x and for every y , $x + y > 10$ ”

Let $P(x, y)$ be the statement $x + y > 10$ where the universe of discourse for x, y is the set of integers.

Universal Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “for every x and for every y , $x + y > 10$ ”

Let $P(x, y)$ be the statement $x + y > 10$ where the universe of discourse for x, y is the set of integers.

Answer:

$$\forall x \forall y P(x, y)$$

Universal Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “for every x and for every y , $x + y > 10$ ”

Let $P(x, y)$ be the statement $x + y > 10$ where the universe of discourse for x, y is the set of integers.

Answer:

$$\forall x \forall y P(x, y)$$

Note that we can also use the shorthand

$$\forall x, y P(x, y)$$

Existential Quantifier

Definition

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Definition

The *existential quantification* of a predicate $P(x)$ is the proposition “There exists an x in the universe of discourse such that $P(x)$ is true.” We use the notation

$$\exists x P(x)$$

which can be read “there exists an x ”

Again, if the universe of discourse is finite, $\{n_1, n_2, \dots, n_k\}$, then the existential quantifier is simply the disjunction of all elements:

$$\exists x P(x) \iff P(n_1) \vee P(n_2) \vee \dots \vee P(n_k)$$

Existential Quantifier

Example I

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Let $P(x, y)$ denote the statement, " $x + y = 5$ ".

What does the expression,

$$\exists x \exists y P(x, y)$$

mean?

What universe(s) of discourse make it true?

Existential Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “there exists a real solution to
 $ax^2 + bx - c = 0$ ”

Existential Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “there exists a real solution to $ax^2 + bx - c = 0$ ”

Let $P(x)$ be the statement $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ where the universe of discourse for x is the set of reals. Note here that a, b, c are all fixed constants.

Existential Quantifier

Example II

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “there exists a real solution to $ax^2 + bx - c = 0$ ”

Let $P(x)$ be the statement $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ where the universe of discourse for x is the set of reals. Note here that a, b, c are all fixed constants.

The statement can thus be expressed as

$$\exists x P(x)$$

Existential Quantifier

Example II Continued

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Question: what is the truth value of $\exists xP(x)$?

Question: what is the truth value of $\exists xP(x)$?

Answer: it is false. For any real numbers such that $b^2 < 4ac$, there will only be complex solutions, for these cases no such *real* number x can satisfy the predicate.

How can we make it so that it *is* true?

Question: what is the truth value of $\exists xP(x)$?

Answer: it is false. For any real numbers such that $b^2 < 4ac$, there will only be complex solutions, for these cases no such *real* number x can satisfy the predicate.

How can we make it so that it *is* true?

Answer: change the universe of discourse to the complex numbers, \mathbb{C} .

In general, when are quantified statements true/false?

Statement	True When	False When
$\forall xP(x)$	$P(x)$ is true for every x .	There is an x for which $P(x)$ is false.
$\exists xP(x)$	There is an x for which $P(x)$ is true.	$P(x)$ is false for every x .

Table: Truth Values of Quantifiers

Existential and universal quantifiers can be used together to quantify a predicate statement; for example,

$$\forall x \exists y P(x, y)$$

is perfectly valid. However, you must be careful—it must be read left to right.

For example, $\forall x \exists y P(x, y)$ is not equivalent to $\exists y \forall x P(x, y)$. Thus, ordering is important.

For example:

- $\forall x \exists y \text{Loves}(x, y)$: everybody loves somebody
- $\exists y \forall x \text{Loves}(x, y)$: There is someone loved by everyone

Those expressions do not mean the same thing!

Note that $\exists y \forall x P(x, y) \rightarrow \forall x \exists y P(x, y)$, but the converse does not hold

However, you *can* commute *similar* quantifiers; $\exists x \exists y P(x, y)$ is equivalent to $\exists y \exists x P(x, y)$ (which is why our shorthand was valid).

Mixing Quantifiers

Truth Values

Statement	True When	False When
$\forall x \forall y P(x, y)$	$P(x, y)$ is true for every pair x, y .	There is at least one pair, x, y for which $P(x, y)$ is false.
$\forall x \exists y P(x, y)$	For every x , there is a y for which $P(x, y)$ is true.	There is an x for which $P(x, y)$ is false for every y .
$\exists x \forall y P(x, y)$	There is an x for which $P(x, y)$ is true for every y .	For every x , there is a y for which $P(x, y)$ is false.
$\exists x \exists y P(x, y)$	There is at least one pair x, y for which $P(x, y)$ is true.	$P(x, y)$ is false for every pair x, y .

Table: Truth Values of 2-variate Quantifiers

Mixing Quantifiers

Example I

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express, in predicate logic, the statement that there are an infinite number of integers.

Mixing Quantifiers

Example I

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express, in predicate logic, the statement that there are an infinite number of integers.

Let $P(x, y)$ be the statement that $x < y$. Let the universe of discourse be the integers, \mathbb{Z} .

Mixing Quantifiers

Example I

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express, in predicate logic, the statement that there are an infinite number of integers.

Let $P(x, y)$ be the statement that $x < y$. Let the universe of discourse be the integers, \mathbb{Z} .

Then the statement can be expressed by the following.

$$\forall x \exists y P(x, y)$$

Express the *commutative law of addition* for \mathbb{R} .

Mixing Quantifiers

Example II: More Mathematical Statements

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the *commutative law of addition* for \mathbb{R} .

We want to express that for every pair of reals, x, y the following identity holds:

$$x + y = y + x$$

Express the *commutative law of addition* for \mathbb{R} .

We want to express that for every pair of reals, x, y the following identity holds:

$$x + y = y + x$$

Then we have the following:

$$\forall x \forall y (x + y = y + x)$$

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
FunctionsPropositional
Functions

Quantifiers

Universal
Quantifier
Existential
QuantifierMixing
QuantifiersBinding
Variables
NegationLogic
ProgrammingTranscribing
English into
Logic

Further

Express the *multiplicative inverse law* for (nonzero) rationals
 $\mathbb{Q} \setminus \{0\}$.

Mixing Quantifiers

Example II: More Mathematical Statements Continued

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the *multiplicative inverse law* for (nonzero) rationals $\mathbb{Q} \setminus \{0\}$.

We want to express that for every real number x , there exists a real number y such that $xy = 1$.

Express the *multiplicative inverse law* for (nonzero) rationals $\mathbb{Q} \setminus \{0\}$.

We want to express that for every real number x , there exists a real number y such that $xy = 1$.

Then we have the following:

$$\forall x \exists y (xy = 1)$$

Mixing Quantifiers

Example II: False Mathematical Statements

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Is commutativity for subtraction valid over the reals?

Mixing Quantifiers

Example II: False Mathematical Statements

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Is commutativity for subtraction valid over the reals?

That is, for all pairs of real numbers x, y does the identity $x - y = y - x$ hold? Express this using quantifiers.

Is commutativity for subtraction valid over the reals?

That is, for all pairs of real numbers x, y does the identity $x - y = y - x$ hold? Express this using quantifiers.

The expression is

$$\forall x \forall y (x - y = y - x)$$

Mixing Quantifiers

Example II: False Mathematical Statements Continued

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Is there a multiplicative inverse law over the nonzero integers?

Mixing Quantifiers

Example II: False Mathematical Statements Continued

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier

Existential
Quantifier

Mixing
Quantifiers

Binding
Variables

Negation

Logic
Programming

Transcribing
English into
Logic

Further

Is there a multiplicative inverse law over the nonzero integers?

That is, for every integer x does there exist a y such that
 $xy = 1$?

Mixing Quantifiers

Example II: False Mathematical Statements Continued

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Is there a multiplicative inverse law over the nonzero integers?

That is, for every integer x does there exist a y such that
 $xy = 1$?

This is false, since we can find a *counter example*. Take any integer, say 5 and multiply it with another integer, y . If the statement held, then $5 = 1/y$, but for any (nonzero) integer y ,
 $|1/y| \leq 1$.

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

Mixing Quantifiers

Exercise

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Future

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

- Let $P(x, y)$ be the expression “ $x + y = y$ ”.

Mixing Quantifiers

Exercise

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Further

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

- Let $P(x, y)$ be the expression “ $x + y = y$ ”.
- Let $Q(x, y)$ be the expression “ $xy = x$ ”.

Mixing Quantifiers

Exercise

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Final

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

- Let $P(x, y)$ be the expression “ $x + y = y$ ”.
- Let $Q(x, y)$ be the expression “ $xy = x$ ”.
- Then the expression is

$$\exists x \forall y (P(x, y) \wedge Q(x, y))$$

Mixing Quantifiers

Exercise

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Fall 2013

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

- Let $P(x, y)$ be the expression “ $x + y = y$ ”.
- Let $Q(x, y)$ be the expression “ $xy = x$ ”.
- Then the expression is

$$\exists x \forall y (P(x, y) \wedge Q(x, y))$$

- Over what universe(s) of discourse does this statement hold?

Mixing Quantifiers

Exercise

Predicate
Logic and
Quantifiers

CSE235

Introduction

Propositional
Functions

Propositional
Functions

Quantifiers

Universal
Quantifier
Existential
Quantifier

Mixing
Quantifiers

Binding
Variables
Negation

Logic
Programming

Transcribing
English into
Logic

Final 33

Express the statement “there is a number x such that when it is added to any number, the result is that number, and if it is multiplied by any number, the result is x ” as a logical expression.

Solution:

- Let $P(x, y)$ be the expression “ $x + y = y$ ”.
- Let $Q(x, y)$ be the expression “ $xy = x$ ”.
- Then the expression is

$$\exists x \forall y (P(x, y) \wedge Q(x, y))$$

- Over what universe(s) of discourse does this statement hold?
- This is the *additive identity law* and holds for $\mathbb{N}, \mathbb{Z}, \mathbb{R}, \mathbb{Q}$ but does not hold for \mathbb{Z}^+ .

When a quantifier is used on a variable x , we say that x is *bound*. If no quantifier is used on a variable in a predicate statement, it is called *free*.

Example

In the expression $\exists x \forall y P(x, y)$ both x and y are bound.
In the expression $\forall x P(x, y)$, x is bound, but y is free.

A statement is called a *well-formed formula*, when all variables are properly quantified.

The set of all variables bound by a common quantifier is the *scope* of that quantifier.

Example

In the expression $\exists x, y \forall z P(x, y, z, c)$ the scope of the existential quantifier is $\{x, y\}$, the scope of the universal quantifier is just z and c has no scope since it is free.

Just as we can use negation with propositions, we can use them with quantified expressions.

Lemma

Let $P(x)$ be a predicate. Then the following hold.

$$\neg\forall xP(x) \equiv \exists x\neg P(x)$$

$$\neg\exists xP(x) \equiv \forall x\neg P(x)$$

This is essentially a quantified version of De Morgan's Law (in fact if the universe of discourse is finite, it is *exactly* De Morgan's law).

Statement	True When	False When
$\neg\exists xP(x) \equiv \forall x\neg P(x)$	For every x , $P(x)$ is false.	There is an x for which $P(x)$ is true.
$\neg\forall xP(x) \equiv \exists x\neg P(x)$	There is an x for which $P(x)$ is false.	$P(x)$ is true for every x .

Table: Truth Values of Negated Quantifiers

Prolog (Programming in Logic) is a programming language based on (a restricted form of) Predicate Calculus. It was developed by the logicians of the artificial intelligence community for symbolic reasoning.

- Prolog allows the user to express facts and rules
- Facts are propositional functions: `student(juana)`, `enrolled(juana,cse235)`, `instructor(patel,cse235)`, etc.

- Rules are implications with conjunctions:
`teaches(X,Y) :- instructor(X,Z), enrolled(Y,Z)`

- Prolog answers queries such as:

`?enrolled(juana,cse478)`

`?enrolled(X,cse478)`

`?teaches(X,juana)`

by binding variables and doing theorem proving (i.e., applying inference rules) as we will see in Section 1.5.

Logic is more precise than English.

Transcribing English to Logic and vice versa can be tricky.

When writing statements with quantifiers, *usually* the correct meaning is conveyed with the following combinations:

- Use \forall with \Rightarrow

Example: $\forall x \text{Lion}(x) \Rightarrow \text{Fierce}(x)$

$\forall x \text{Lion}(x) \wedge \text{Fierce}(x)$ means “everyone is a lion and everyone is fierce”

- Use \exists with \wedge

Example: $\exists x \text{Lion}(x) \wedge \text{Drinks}(x, \text{coffee})$: holds when you have at least one lion that drinks coffee

$\exists x \text{Lion}(x) \Rightarrow \text{Drinks}(x, \text{coffee})$ holds when you have people even though no lion drinks coffee.

Examples? Exercises?

- Rewrite the expression,
$$\neg \forall x (\exists y \forall z P(x, y, z) \wedge \exists z \forall y P(x, y, z))$$
- Let $P(x, y)$ denote “ x is a factor of y ” where $x \in \{1, 2, 3, \dots\}$ and $y \in \{2, 3, 4, \dots\}$. Let $Q(y)$ denote “ $\forall x [P(x, y) \rightarrow ((x = y) \vee (x = 1))]$ ”. When is $Q(y)$ true?

Examples? Exercises?

- Rewrite the expression,
 $\neg \forall x (\exists y \forall z P(x, y, z) \wedge \exists z \forall y P(x, y, z))$
- Answer: Use the negated quantifiers and De Morgan's law.

$$\exists x (\forall y \exists z \neg P(x, y, z) \vee \forall z \exists y \neg P(x, y, z))$$

- Let $P(x, y)$ denote “ x is a factor of y ” where $x \in \{1, 2, 3, \dots\}$ and $y \in \{2, 3, 4, \dots\}$. Let $Q(y)$ denote “ $\forall x [P(x, y) \rightarrow ((x = y) \vee (x = 1))]$ ”. When is $Q(y)$ true?

Examples? Exercises?

- Rewrite the expression,
$$\neg \forall x (\exists y \forall z P(x, y, z) \wedge \exists z \forall y P(x, y, z))$$
- Answer: Use the negated quantifiers and De Morgan's law.

$$\exists x (\forall y \exists z \neg P(x, y, z) \vee \forall z \exists y \neg P(x, y, z))$$

- Let $P(x, y)$ denote “ x is a factor of y ” where $x \in \{1, 2, 3, \dots\}$ and $y \in \{2, 3, 4, \dots\}$. Let $Q(y)$ denote “ $\forall x [P(x, y) \rightarrow ((x = y) \vee (x = 1))]$ ”. When is $Q(y)$ true?
- Answer: Only when y is a prime number.

Some students wondered if

$$\forall x, y P(x, y) \equiv \forall x P(x, y) \wedge \forall y P(x, y)$$

This is certainly not true. In the left-hand side, both x and y are bound. In the right-hand side, x is bound in the first predicate, but y is free. In the second predicate, y is bound but x is free.

All variables that occur in a propositional function must be bound to turn it into a proposition.

Thus, the left-hand side is a proposition, but the right-hand side is not. How can they be equivalent?