

Sequences & Summations

Slides by Christopher M. Bourke
Instructor: Berthe Y. Choueiry

Fall 2007

Computer Science & Engineering 235
Introduction to Discrete Mathematics
Section 2.4 of Rosen
cse235@cse.unl.edu

Notes

Sequences & Summations

Though you should be (at least intuitively) familiar with sequences and summations, we give a quick review.

Notes

Sequences

Definition

A *sequence* is a function from a subset of integers to a set S . We use the notation(s):

$$\{a_n\} \quad \{a_n\}_n^\infty \quad \{a_n\}_{n=0}^\infty \quad \{a_n\}_{n=0}$$

Each a_n is called the n -th *term* of the sequence.

We rely on context to distinguish between a sequence and a set; though there is a connection.

Notes

Sequences

Example

Example

Consider the sequence

$$\left\{ \left(1 + \frac{1}{n} \right)^n \right\}_{n=1}^{\infty}$$

The terms are

$$a_1 = (1 + 1)^1 = 2.00000$$

$$a_2 = \left(1 + \frac{1}{2} \right)^2 = 2.25000$$

$$a_3 = \left(1 + \frac{1}{3} \right)^3 = 2.37037$$

$$a_4 = \left(1 + \frac{1}{4} \right)^4 = 2.44140$$

$$a_5 = \left(1 + \frac{1}{5} \right)^5 = 2.48832$$

What is this sequence?

Notes

Sequences

Example

The sequence corresponds to e :

$$\lim_{n \rightarrow \infty} \left\{ \left(1 + \frac{1}{n} \right)^n \right\} = e = 2.71828 \dots$$

Notes

Sequences

Example II

Example

The sequence

$$\{h_n\}_{n=1}^{\infty} = \frac{1}{n}$$

is known as the *harmonic sequence*.

The sequence is simply

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots$$

This sequence is particularly interesting because its summation is *divergent*;

$$\sum_{n=1}^{\infty} \frac{1}{n} = \infty$$

Notes

Progressions I

Definition

A *geometric progression* is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

Where $a \in \mathbb{R}$ is called the *initial term* and $r \in \mathbb{R}$ is the *common ratio*.

A geometric progression is a *discrete* analogue of the exponential function

$$f(x) = ar^x$$

Notes

Progressions II

Definition

An *arithmetic progression* is a sequence of the form

$$a, a + d, a + 2d, a + 3d, \dots, a + nd, \dots$$

Where $a \in \mathbb{R}$ is called the *initial term* and $r \in \mathbb{R}$ is the *common difference*.

Again, an arithmetic progression is a discrete analogue of the linear function,

$$f(x) = dx + a$$

Notes

Progressions III

Example

A common geometric progression in computer science is

$$\{a_n\} = \frac{1}{2^n}$$

Here, $a = 1$ and $r = \frac{1}{2}$

Table 1 on Page 153 (Rosen) has useful sequences.

Notes

Summations I

You should be very familiar with Summation notation:

$$\sum_{j=m}^n a_j = a_m + a_{m+1} + \cdots + a_{n-1} + a_n$$

Here, j is the *index of summation*, m is the *lower limit*, and n is the *upper limit*.

Often times, it is useful to change the lower/upper limits; which can be done in a straightforward manner (though we must be careful).

$$\sum_{j=1}^n a_j = \sum_{j=0}^{n-1} a_{j+1}$$

Sometimes we can express a summation in *closed form*. Geometric series, for example:

Summations II

Theorem

For $a, r \in \mathbb{R}$, $r \neq 0$,

$$\sum_{i=0}^n ar^i = \begin{cases} \frac{ar^{n+1}-a}{r-1} & \text{if } r \neq 1 \\ (n+1)a & \text{if } r = 1 \end{cases}$$

Summations III

Double summations often arise when analyzing an algorithm.

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^i a_j &= a_1 + \\ &\quad a_1 + a_2 + \\ &\quad a_1 + a_2 + a_3 + \\ &\quad \cdots \\ &\quad a_1 + a_2 + a_3 + \cdots + a_n \end{aligned}$$

Summations can also be indexed over *elements in a set*.

$$\sum_{s \in S} f(s)$$

Table 2 on Page 157 (Rosen) has useful summations.

Notes

Notes

Notes

Series

When we take the sum of a sequence, we get a *series*. We've already seen a closed form for geometric series.

Some other useful closed forms include the following.

$$\sum_{i=l}^u 1 = u - l + 1, \text{ for } l \leq u$$
$$\sum_{i=0}^n i = \frac{n(n+1)}{2}$$
$$\sum_{i=0}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$
$$\sum_{i=0}^n i^k \approx \frac{1}{k+1} n^{k+1}$$

Notes

Infinite Series I

Though we will mostly deal with finite series (i.e. an upper limit of n for a fixed integer), *infinite series* are also useful.

Example

Notes

Infinite Series II

Consider the geometric series

$$\sum_{n=0}^{\infty} \frac{1}{2^n} = 1 + \frac{1}{2} + \frac{1}{4} + \dots$$

This series converges to 2. However, the geometric series

$$\sum_{n=0}^{\infty} 2^n = 1 + 2 + 4 + 8 + \dots$$

does not converge. However, note that $\sum_{n=0}^n 2^n = 2^{n+1} - 1$

In fact, we can generalize this as follows.

Lemma

Notes

Infinite Series III

A geometric series converges if and only if the absolute value of the common ratio is less than 1.

Notes
