

Partial Orders

Slides by Christopher M. Bourke
Instructor: Berthe Y. Choueiry

Spring 2006

Computer Science & Engineering 235
Introduction to Discrete Mathematics
Sections 7.6 of Rosen
cse235@cse.unl.edu

Notes

Partial Orders I

Motivating Introduction

Consider the recent renovation of Avery Hall. In this process several things had to be done.

- ▶ Remove Asbestos
- ▶ Replace Windows
- ▶ Paint Walls
- ▶ Refinish Floors
- ▶ Assign Offices
- ▶ Move in Office-Furniture.

Notes

Partial Orders II

Motivating Introduction

Clearly, some things had to be done before others could even begin—Asbestos had to be removed before *anything*; painting had to be done before the floors to avoid ruining them, etc.

On the other hand, several things could have been done concurrently—painting could be done while replacing the windows and assigning office could have been done at anytime.

Such a scenario can be nicely modeled using *partial orderings*.

Notes

Partial Orderings I

Definition

Definition

A relation R on a set S is called a *partial order* if it is reflexive, antisymmetric and transitive. A set S together with a partial ordering R is called a *partially ordered set* or *poset* for short and is denoted

$$(S, R)$$

Partial orderings are used to give an order to sets that may not have a natural one. In our renovation example, we could define an ordering such that $(a, b) \in R$ if a *must* be done before b can be done.

Notes

Partial Orderings II

Definition

We use the notation

$$a \preceq b$$

to indicate that $(a, b) \in R$ is a partial order and

$$a \prec b$$

when $a \neq b$.

The notation \prec is not to be mistaken for "less than equal to." Rather, \prec is used to denote *any* partial ordering.

Latex notation: `\preccurlyeq`, `\prec`.

Notes

Comparability

Definition

The elements a and b of a poset (S, \preceq) are called *comparable* if either $a \preceq b$ or $b \preceq a$. When $a, b \in S$ such that neither are comparable, we say that they are *incomparable*.

Looking back at our renovation example, we can see that

$$\text{Remove Asbestos} \prec a_i$$

for all activities a_i . Also,

$$\text{Paint Walls} \prec \text{Refinish Floors}$$

Some items are also incomparable—replacing windows can be done before, after or during the assignment of offices.

Notes

Total Orders

Definition

If (S, \preceq) is a poset and every two elements of S are comparable, S is called a *totally ordered set*. The relation \preceq is said to be a *total order*.

Example

The set of integers over the relation “less than equal to” is a total order; (\mathbb{Z}, \leq) since for every $a, b \in \mathbb{Z}$, it must be the case that $a \leq b$ or $b \leq a$.

What happens if we replace \leq with $<$?

Notes

Well-Orderings

Definition

(S, \preceq) is a *well-ordered set* if it is a poset such that \preceq is a total ordering and such that every nonempty subset of S has a *least element*.

Example

The natural numbers along with \leq , (\mathbb{N}, \leq) is a well-ordered set since any subset of \mathbb{N} will have a least element and \leq is a total ordering on \mathbb{N} as before.

However, (\mathbb{Z}, \leq) is not a well-ordered set. Why? Is it totally ordered?

Notes

Principle of Well-Ordered Induction

Well-ordered sets are the basis of the proof technique known as *induction* (more when we cover Chapter 3).

Theorem (Principle of Well-Ordered Induction)

Suppose that S is a well ordered set. Then $P(x)$ is true for all $x \in S$ if

Basis Step: $P(x_0)$ is true for the least element of S and

Induction Step: For every $y \in S$ if $P(x)$ is true for all $x \prec y$ then $P(y)$ is true.

Notes

Principle of Well-Ordered Induction

Proof

Suppose it is not the case that $(P(x))$ holds for all $x \in S \Rightarrow \exists y P(y)$ is false $\Rightarrow A = \{x \in S | P(x) \text{ is false}\}$ is not empty.

Since S is well ordered, A has a least element a .

$P(x_0)$ is true $\Rightarrow a \neq x_0$.

$P(x)$ holds for all $x \in S$ and $x \prec a$, then $P(a)$ holds, by the induction step.

This yields a contradiction. \square

Notes

Lexicographic Orderings I

Lexicographic ordering is the same as any dictionary or phone book—we use alphabetical order starting with the first character in the string, then the next character (if the first was equal) etc. (you can consider “no character” for shorter words to be less than “a”).

Notes

Lexicographic Orderings II

Formally, lexicographic ordering is defined by combining two other orderings.

Definition

Let (A_1, \preceq_1) and (A_2, \preceq_2) be two posets. The *lexicographic ordering* \preceq on the Cartesian product $A_1 \times A_2$ is defined by

$$(a_1, a_2) \preceq (a'_1, a'_2)$$

if $a_1 \prec_1 a'_1$ or if $a_1 = a'_1$ and $a_2 \preceq_2 a'_2$.

Notes

Lexicographic Orderings III

Lexicographic ordering generalizes to the Cartesian product of n sets in the natural way.

Define \preceq on $A_1 \times A_2 \times \cdots \times A_n$ by

$$(a_1, a_2, \dots, a_n) \prec (b_1, b_2, \dots, b_n)$$

if $a_1 \prec b_1$ or if there is an integer $i > 0$ such that

$$a_1 = b_1, a_2 = b_2, \dots, a_i = b_i$$

and $a_{i+1} \prec b_{i+1}$

Notes

Lexicographic Orderings I

Strings

Consider the two non-equal strings $a_1a_2 \cdots a_m$ and $b_1b_2 \cdots b_n$ on a poset S .

Let $t = \min(n, m)$ and \prec is the lexicographic ordering on S^t .

$a_1a_2 \cdots a_m$ is less than $b_1b_2 \cdots b_n$ if and only if

- ▶ $(a_1, a_2, \dots, a_t) \prec (b_1, b_2, \dots, b_t)$, or
- ▶ $(a_1, a_2, \dots, a_t) = (b_1, b_2, \dots, b_t)$ and $m < n$

Notes

Hasse Diagrams

As with relations and functions, there is a convenient graphical representation for partial orders—*Hasse Diagrams*.

Consider the digraph representation of a partial order—since we *know* we are dealing with a partial order, we *implicitly* know that the relation must be reflexive and transitive. Thus we can simplify the graph as follows:


- ▶ Remove all self-loops.
- ▶ Remove all transitive edges.
- ▶ Make the graph direction-less—that is, we can assume that the orientations are *upwards*.

The resulting diagram is far simpler.

Notes

Hasse Diagram

Example




Remove Self-Loops

Notes

Hasse Diagram

Example




Remove Transitive Loops

Notes

Hasse Diagram

Example




Remove Orientation

Notes

Hasse Diagram

Example



Hasse Diagram!

Notes

Hasse Diagrams

Example

Of course, you need not always start with the complete relation in the partial order and then trim everything. Rather, you can build a Hasse directly from the partial order.

Example

Draw a Hasse diagram for the partial ordering


$$\{(a, b) \mid a \mid b\}$$

on $\{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$ (these are the divisors of 60 which form the basis of the ancient Babylonian base-60 numeral system)

Notes

Hasse Diagrams

Example Answer



Notes

Extremal Elements I

Summary

We will define the following terms:

- ▶ A maximal/minimal element in a poset (S, \preceq) .
- ▶ The maximum (greatest)/minimum (least) element of a poset (S, \preceq) .
- ▶ An upper/lower bound element of a subset A of a poset (S, \preceq) .
- ▶ The greatest upper/least lower bound element of a subset A of a poset (S, \preceq) .
- ▶ Lattice

Notes

Extremal Elements I

Definition

An element a in a poset (S, \preceq) is called *maximal* if it is not less than any other element in S . That is,

$$\nexists b \in S (a \prec b)$$

If there is one *unique* maximal element a , we call it the *maximum* element (or the *greatest element*).

Notes

Extremal Elements II

Definition

An element a in a poset (S, \preceq) is called *minimal* if it is not greater than any other element in S . That is,

$$\nexists b \in S (b \prec a)$$

If there is one *unique* minimal element a , we call it the *minimum* element (or the *least element*).

Notes

Extremal Elements III

Definition

Let (S, \preceq) be a poset and let $A \subseteq S$. If u is an element of S such that $a \preceq u$ for all elements $a \in A$ then u is an *upper bound* of A .

An element x that is an upper bound on a subset A and is less than all other upper bounds on A is called the *least upper bound* on A . We abbreviate "lub".

Notes

Extremal Elements IV

Definition

Let (S, \preceq) be a poset and let $A \subseteq S$. If l is an element of S such that $l \preceq a$ for all elements $a \in A$ then l is a *lower bound* of A .


An element x that is a lower bound on a subset A and is greater than all other lower bounds on A is called the *greatest lower bound* on A . We abbreviate "glb".

Notes

Extremal Elements

Example I

Example



What are the minimal, maximal, minimum, maximum elements?


- ▶ Minimal: $\{a, b\}$
- ▶ Maximal: $\{c, d\}$
- ▶ There are no unique minimal or maximal elements.

Notes

Extremal Elements

Example II

Example



What are the lower/upper bounds and glb/lub of the sets $\{d, e, f\}$, $\{a, c\}$ and $\{b, d\}$

Notes

Extremal Elements

Example II

$\{d, e, f\}$

- ▶ Lower Bounds: \emptyset , thus no glb either.
- ▶ Upper Bounds: \emptyset , thus no lub either.

$\{a, c\}$

- ▶ Lower Bounds: \emptyset , thus no glb either.
- ▶ Upper Bounds: $\{h\}$, since its unique, lub is also h .

$\{b, d\}$


- ▶ Lower Bounds: $\{b\}$ and so also glb.
- ▶ Upper Bounds: $\{d, g\}$ and since $d \prec g$, the lub is d .

Notes

Extremal Elements

Example III

Example



Minimal/Maximal elements?

- ▶ Minimal & Minimum Element: a .
- ▶ Maximal Elements: b, d, i, j .

Bounds, glb, lub of $\{c, e\}$?

- ▶ Lower Bounds: $\{a, c\}$, thus glb is c .
- ▶ Upper Bounds: $\{e, f, g, h, i, j\}$ thus lub is e

Bounds, glb, lub of $\{b, i\}$?

- ▶ Lower Bounds: $\{a\}$, thus glb is a .
- ▶ Upper Bounds: \emptyset , thus lub DNE.

Notes

Lattices

A special structure arises when every pair of elements in a poset has a lub and glb.

Definition


A partially ordered set in which every pair of elements has both a least upper bound and a greatest lower bound is called a *lattice*.

Notes

Lattices

Example

Is the example from before a lattice?




No, since the pair (b, c) do not have a least upper bound.

Notes

Lattices

Example

What if we modified it as follows?



Yes, it is now a lattice, since for any pair, there is a lub & glb.

Notes

Lattices

To show that a partial order is not a lattice, it suffices to find a pair that does not have a lub/glb.

For a pair not to have a lub/glb, they must first be *incomparable*. (Why?)

You can then view the upper/lower bounds on a pair as a sub-hasse diagram; if there is no *minimum* element in this sub-diagram, then it is not a lattice.

Notes

Topological Sorting

Introduction

Let us return to the introductory example of the Avery renovation. Now that we have got a partial order model, it would be nice to actually create a concrete schedule.

That is, given a partial order, we would like to transform it into a *total order* that is *compatible* with the partial order.

A total order is compatible if it doesn't violate any of the original relations in the partial ordering.

Essentially, we are simply imposing an order on incomparable elements in the partial order.

Notes

Preliminaries

Before we give the algorithm, we need some tools to justify its correctness.

Fact

Every finite, nonempty poset (S, \preceq) has a minimal element.

We will prove by a form of *reductio ad absurdum*.

Notes

Preliminaries

Proof

Proof.

Assume to the contrary that a nonempty, finite (WLOG, assume $|S| = n$) poset (S, \preceq) has no minimal element. In particular, a_1 is not a minimal element.

If a_1 is not minimal, then there exists a_2 such that $a_2 \prec a_1$. But also, a_2 is not minimal by the assumption.

Therefore, there exists a_3 such that $a_3 \prec a_2$. This process proceeds until we have the last element, a_n thus,

$$a_n \prec a_{n-1} \prec \cdots a_2 \prec a_1$$

thus by definition a_n is the minimal element. \square

Notes

Topological Sorting

Intuition

The idea to topological sorting is that we start with a poset (S, \preceq) and remove a minimal element (choosing arbitrarily if there are more than one). Such an element is guaranteed to exist by the previous fact.

As we remove each minimal element, the set shrinks. Thus, we are guaranteed the algorithm will halt in a finite number of steps.

Furthermore, the order in which elements are removed is a total order;

$$a_1 \prec a_2 \prec \cdots \prec a_n$$

We now present the algorithm itself.

Notes

Topological Sorting

Algorithm

TOPOLOGICAL SORT

```
INPUT      :  $(S, \preceq)$  a poset with  $|S| = n$ 
OUTPUT     : A total ordering  $(a_1, a_2, \dots, a_n)$ 
1  $k = 1$ 
2 WHILE  $S \neq \emptyset$  DO
3    $a_k \leftarrow$  a minimal element in  $S$ 
4    $S = S \setminus \{a_k\}$ 
5    $k = k + 1$ 
6 END
7 return  $(a_1, a_2, \dots, a_n)$ 
```


Notes

Topological Sorting

Example

Example

Find a compatible ordering (topological ordering) of the poset represented by the diagram below.



Notes

Conclusion

Questions?

Notes
