### **Sequences & Summations**

#### **Section 2.4 of Rosen**

Spring 2011

**CSCE 235 Introduction to Discrete Structures** 

Course web-page: cse.unl.edu/~cse235

Questions: cse235@cse.unl.edu

#### **Outline**

Although you are (more or less) familiar with sequences and summations, we give a quick review

- Sequences
  - Definition, 2 examples
- Progressions: Special sequences
  - Geometric, arithmetic
- Summations
  - Careful when changing lower/upper limits
- Series: Sum of the elements of a sequence
  - Examples, infinite series, convergence of a geometric series

#### Sequences

• **Definition**: A <u>sequence</u> is a function from a subset of integers to a set S. We use the notation(s):

$$\{a_n\}$$
  $\{a_n\}_n^{\infty}$   $\{a_n\}_{n=0}^{\infty}$ 

- Each a<sub>n</sub> is called the n<sup>th</sup> term of the sequence
- We rely on the context to distinguish between a sequence and a set, although they are distinct structures

## Sequences: Example 1

Consider the sequence

$$\{(1 + 1/n)^n\}_{n=1}^{\infty}$$

The terms of the sequence are:

$$a_1 = (1 + 1/1)^1 = 2.00000$$
 $a_2 = (1 + 1/2)^2 = 2.25000$ 
 $a_3 = (1 + 1/3)^3 = 2.37037$ 
 $a_4 = (1 + 1/4)^4 = 2.44140$ 
 $a_5 = (1 + 1/5)^5 = 2.48832$ 

- What is this sequence?
- The sequence corresponds to  $\lim_{n\to\infty} \{(1+1/n)^n\}_{n=1}^n = e = 2.71828..$

# Sequences: Example 2

- The sequence:  $\{h_n\}_{n=1}^{\infty} = 1/n$ is known as the <u>harmonic</u> sequence
- The sequence is simply:

 This sequence is particularly interesting because its summation is divergent:

$$\sum_{n=1}^{\infty} (1/n) = \infty$$

### Progressions: Geometric

 Definition: A geometric progression is a sequence of the form

#### Where:

- a∈R is called the <u>initial term</u>
- r∈R is called the <u>common ratio</u>
- A geometric progression is a <u>discrete</u> analogue of the exponential function

$$f(x) = ar^x$$

## Geometric Progressions: Examples

 A common geometric progression in Computer Science is:

$${a_n} = 1/2^n$$

with a=1 and r=1/2

- Give the initial term and the common ratio of
  - $-\{b_n\}$  with  $b_n = (-1)^n$
  - $\{c_n\}$  with  $c_n = 2(5)^n$
  - $\{d_n\}$  with  $d_n = 6(1/3)^n$

## Progressions: Arithmetic

 Definition: An <u>arithmetric progression</u> is a sequence of the form

#### Where:

- $-a \in R$  is called the <u>initial term</u>
- $-d \in R$  is called the common difference
- An arithmetic progression is a <u>discrete</u> analogue of the linear function

$$f(x) = dx+a$$

# Arithmetic Progressions: Examples

- Give the initial term and the common difference of
  - $-\{s_n\}$  with  $s_n = -1 + 4n$
  - $-\{t_n\}$  with  $s_n = 7 3n$

### More Examples

 Table 1 on Page 153 (Rosen) has some useful sequences:

$$\{n^2\}_{n=1}^{\infty}$$
,  $\{n^3\}_{n=1}^{\infty}$ ,  $\{n^4\}_{n=1}^{\infty}$ ,  $\{2^n\}_{n=1}^{\infty}$ ,  $\{3^n\}_{n=1}^{\infty}$ ,  $\{n!\}_{n=1}^{\infty}$ 

#### **Outline**

# Although you are (more or less) familiar with sequences and summations, we give a quick review

- Sequences
  - Definition, 2 examples
- Progressions: Special sequences
  - Geometric, arithmetic
- Summations
  - Careful when changing lower/upper limits
- Series: Sum of the elements of a sequence
  - Examples, infinite series, convergence of a geometric series

# Summations (1)

You should be by now familiar with the summation notation:

$$\Sigma_{j=m}^{n} (a_j) = a_m + a_{m+1} + ... + a_{n-1} + a_n$$

#### Here

- j is the index of the summation
- m is the lower limit
- n is the upper limit
- Often times, it is useful to change the lower/upper limits, which can be done in a straightforward manner (although we must be very careful):

$$\Sigma_{i=1}^{n} (a_i) = \Sigma_{i=0}^{n-1} (a_{i+1})$$

# Summations (2)

- Sometimes we can express a summation in closed form, as for geometric series
- **Theorem**: For a,  $r \in \mathbb{R}$ ,  $r \neq 0$

$$\Sigma_{i=0}^{n} (ar^{i}) = \begin{cases} (ar^{n+1}-a)/(r-1) & \text{if } r \neq 1 \\ (n+1)a & \text{if } r = 1 \end{cases}$$

 Closed form = analytical expression using a bounded number of well-known functions, does not involved an infinite series or use of recursion

# Summations (3)

Double summations often arise when analyzing an algorithm

$$\Sigma_{i=1}^{n} \Sigma_{j=1}^{i}(a_{j}) = a_{1} + a_{1} + a_{2} + a_{3} + a_{2} + a_{3} + a_{2} + a_{3} + a_{1} + a_{2} + a_{3} + a_{2} + a_{3} + a_{3} + a_{4} + a_{4} + a_{5} + a_{$$

Summations can also be indexed over elements in a set:

$$\Sigma_{s \in S} f(s)$$

Table 2 on Page 157 (Rosen) has very useful summations.
 Exercises 2.4.14—18 are great material to practice on.

#### **Outline**

Although you are (more or less) familiar with sequences and summations, we give a quick review

- Sequences
  - Definition, 2 examples
- Progressions: Special sequences
  - Geometric, arithmetic
- Summations
  - Careful when changing lower/upper limits
- Series: Sum of the elements of a sequence
  - Examples, infinite series, convergence of a geometric series

#### Series

- When we take the <u>sum of a sequence</u>, we get a <u>series</u>
- We have already seen a closed form for geometric series
- Some other useful closed forms include the following:

$$-\sum_{i=k}^{u} 1 = u-k+1$$
, for k≤u

$$-\sum_{i=0}^{n} i = n(n+1)/2$$

$$-\sum_{i=0}^{n} (i^2) = n(n+1)(2n+1)/6$$

$$-\sum_{i=0}^{n} (i^{k}) \approx n^{k+1}/(k+1)$$

#### Infinite Series

- Although we will mostly deal with finite series (i.e., an upper limit of n for fixed integer), inifinite series are also useful
- Consider the following geometric series:

$$-\sum_{n=0}^{\infty} (1/2^n) = 1 + 1/2 + 1/4 + 1/8 + \dots$$
 converges to 2

$$-\sum_{n=0}^{\infty} (2^n) = 1 + 2 + 4 + 8 + \dots$$
 does not converge

• However note:  $\Sigma_{n=0}^{n} (2^{n}) = 2^{n+1} - 1 (a=1,r=2)$ 

#### Infinite Series: Geometric Series

- In fact, we can generalize that fact as follows
- Lemma: A geometric series converges <u>if and</u> only <u>if</u> the absolute value of the common ratio is less than 1

```
When |r| < 1,

\lim_{n \to \infty} \sum_{i=0}^{n} (ar^{i}) = \lim_{n \to \infty} \sum_{i=0}^{n} (ar^{n+1}-a)/(r-1)

= a/(1-r)
```